



# Bodleian Libraries

UNIVERSITY OF OXFORD


This book is part of the collection held by the Bodleian Libraries and scanned by Google, Inc. for the Google Books Library Project.

For more information see:

<http://www.bodleian.ox.ac.uk/dbooks>



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 2.0 UK: England & Wales (CC BY-NC-SA 2.0) licence.




G. 86. a. 13



E. BIBL. RADCL.

~~H. 51/3~~  
13.

~~G. 4/4~~ ~~71 A. 13.~~  
13



CR. 5.90/13 C

18933 d. 215/13

J-48 col:

March 1

April 1

May 1

June 1

July 1


Aug 1





TO  
THE ROYAL COLLEGE  
OF  
PHYSICIANS OF LONDON,  
AS A TRIBUTE OF RESPECT  
TO  
THAT LEARNED BODY,  
*THIS THIRTEENTH VOLUME*  
OF THE  
NATURALIST'S MISCELLANY  
IS INSCRIBED  
BY  
GEORGE SHAW,  
E. R. NODDER.





COLLEGII REGALIS,  
MEDICORUM LONDINENSIIUM,

PRÆSIDI

SOCIISQUE ORNATISSIMIS,

DECIMUM TERTIUM

HUNC

NATURÆ VIVARIII

*FASCICULUM,*

D. D. D.

GEORGIUS SHAW,

E. R. NODDER.



## PICUS PILEATUS.



### CHARACTER GENERICUS.

*Rostrum* polyedrum, rectum, apice cuneato.

*Nares* pennis setaceis recumbentibus obtectæ.

*Lingua* teres, lumbriciformis, longissima, mucronata, apice retrorsum aculeata setis.

*Pedes* scansorii.

### CHARACTER SPECIFICUS, &c.

PICUS niger, crista rubra, temporibus alisque albis maculis.

*Lin. Syst. Nat.* p. 173.

PICUS Virginianus PILEATUS.

*Briss. av.* 4. p. 29.

---

Pico qui *principalis* appellatur seu *albirostris*, affinis admodum est picus pileatus; paulo tamen minor, alium quoque habet colorem rostri. In America septentrionali generatus amat præcipue Carolinam et Virginiam. Arbores grandiores terebrando excavare, triticoque Indico, quod *Maiz* dicitur, non leve damnum solet inferre. Magnitudine fere corvum vulgarem æquat.

## PILEATED WOODPECKER.



### GENERIC CHARACTER.

*Bill* polyedral, strait, wedge-shaped towards the tip.

*Nostrils* covered by recumbent setaceous feathers.

*Tongue* worm-shaped, very long, sharp-pointed and edged towards the tip with reversed bristles.

*Feet* scansorial.

### SPECIFIC CHARACTER, &c.

Black WOODPECKER, with red crest, and the temples and wings marked by white spots.

Larger crested WOODPECKER.

*Catesb. Carol.* 1. pl. 17.

PILEATED WOODPECKER.

*Penn. Arct. Zool.* p. 269.

---

This species is very nearly allied to the *Picus principalis* or White-billed Woodpecker, but is of rather smaller size, and has a bill of a different colour. It is a native of North America, and is particularly



A. N.

London Published Sept. 1780 by Elizabeth Nodder



particularly found in Carolina and Virginia, where it forms excavations in the larger trees, and is also said to make considerable havock among the plantations of maize or Indian corn. In size it is not much inferior to a Crow.



## SCOMBER RUBER.



### CHARACTER GENERICUS.

*Caput compressum, læve.*

*Membrana branchiostega radiis septem.*

*Corpus læve, linea laterali postice carinatum.*

*Pinnæ spuriae sæpius versus caudam.*

*Lin. Syst. Nat. p. 492.*

### CHARACTER SPECIFICUS, &c.

SCOMBER corpore rubro.

*Bloch. ichth. 10. p. 63. t. 342.*


Die Rothe Mackrele.

*Bloch. ichth. 10. p. 63.*

---

Colore solo ab omnibus hujus generis facillime distinguitur Scomber ruber. In mari Americano præcipue conspicitur, magnitudine, ut plurimum, sesquipedali.





*Sea bream*

585

494

THE  
RED MACKREL.



*GENERIC CHARACTER.*

*Head* compressed, smooth.

*Gill-Membrane* furnished with seven rays.

*Body* smooth, carinated at the hind part by the lateral line.

*Spurious pinnules* (in many species) towards the tail.

*SPECIFIC CHARACTER.*

**MACKREL** with red body.

---

This species is readily distinguished by its colour alone from all its congeners. It is chiefly found in the American seas, and grows to the length of about a foot and half.

CONUS AMMIRALIS.

VAR.

AMM: SUMMUS.

ET

CEDO NULLI.



CHARACTER GENERICUS.

*Animal* Limax.

*Testa* univalvis, convoluta, turbinata.

*Apertura* effusa, longitudinalis, linearis, edentula, basi integra.

*Columella* lævis.

*Lin. Syst. Nat.* p. 1166.

CHARACTER SPECIFICUS, &c.

CONUS testa albo ferrugineoque varia, basi punctato-scabra.

CONUS AMMIRALIS. C. testa basi punctato-scabra.

*Lin. Syst. Nat.* p. 1167.

*Var.* AMMIRALIS summus. C. testa ferruginea maculis albis sparsis, fasciisque quatuor flavis tenuissime reticulatis, tertia cingulo albo maculato.

*Lin. Syst. Nat.* p. 1167.

*Var.*

*Var. Cedo nulli. C. testa testacea albo maculata, cingulisque tribus, supremo composito, omnibus punctatis.*

*Lin. Syst. Nat. p. 1167.*

---

In tot varietates sparguntur testæ quas complectitur *Coni* genus, ut uniuscujusque speciem pro certo definire difficillimum sit: præsertim cum ob scelertas mercatorum fraudes interdum fere impossibile sit falsas et adulterinas a veris et naturalibus discernere.

Ex elegantissimis conorum est ille qui *Ammiralis* dicitur, cujus multæ sunt varietates, quarum princeps est *Ammiralis summus*, in tabulæ nostræ parte superiori depictus. Ex omni autem concharum familia, nulla est ab iis magis deamata, qui in colligendis rarioribus immodicas profundunt pecunias, quam quæ nomine *Cedo nulli* insignita significanter a Linnæo vocatur “pretiosissimus artis perditæ luxus.” Huic proprium et peculiare est quod zonæ ferruginæ seu aurantiæ ordinibus multis macularum parvularum et albarum cingantur margaritis non longe absimilium. Nullum extitisse dicitur specimen pulchrius et perfectius quam quod Hagæ continuere scrinia Domini Lyonetti.

THE  
ADMIRAL CONE.

435

VAR.  
HIGH ADMIRAL.

AND  
CEDO NULLI.



*GENERIC CHARACTER.*

*Animal* resembling a Limax or Slug.

*Shell* more or less conic or pyramidal.

*Aperture* longitudinal, linear, without teeth,  
entire at the base.

*Pillar* smooth.

*SPECIFIC CHARACTER, &c.*

CONE with ferruginous and white variegations,  
and the base roughened with small points.

*Var.* High Admiral. Ferruginous Cone with  
scattered white spots, and four yellow  
bands finely reticulated, the third marked  
by a spotted white line.

*Var.* Cedo nulli. Yellowish-brown Cone spotted  
with white, with three bands, all marked  
by rows of round white specks.

The

465



PN





The varieties which take place in the genus *Conus* are so numerous as to render the precise determination of the species a matter of no easy investigation: it may also be added that in no genus have the arts of dealers been more exerted, nor is it possible to determine whether many of the shells seen in collections are truly natural or not. Among the most elegant of the Cones is the *C. Ammiralis* or *Admiral-Shell*, which admits of many varieties, and of which the chief is the *C. Ammiralis summus* or *High Admiral*, represented on the upper part of the present plate. But of all the tribe that which has most excited the ambition of collectors is the variety termed *Cedo nulli*, which Linnæus emphatically calls “pretiosissimus artis perditæ lux-  
“ us.” This curious shell is distinguished by having the brown or orange-coloured zones ornamented with several rows of small, round, white, pearl-like specks; a particularity not to be found in any of the rest. The most perfect specimen of the *Cedo nulli* is said to have been that in the collection of Mr. Lyonet of the Hague.

## CANCER PHALANGIUM.



### CHARACTER GENERICUS.

*Pedes* octo (rarius sex aut decem); insuper manus duæ chelatae.

*Palpi* sex inæquales.

*Oculi* duo distantes, plurimis pedunculati; elongati, mobiles.

*Mandibula* crassa, cornea.

*Labium* triplex.

*Cauda* articulata, inermis.

*Lin. Syst. Nat. Gmel. p. 2963.*

### CHARACTER SPECIFICUS, &c.

CANCER corpore cordato tuberculato, rostrum bifido, pedibus longissimis gracillimis pilosis.

### CANCER PHALANGIUM.

*Penn. Brit. Zool. 4. p. 7. t. 9. f. 17.*

---

Mira omnino et insolita est huic cancro membrorum longitudo et gracilitas; unde et nomen adeptus est; primo enim visu phalangii seu araneæ speciem potius quam cancrum putares. In oris Europæis generatur, eadem, ut plurimum, magnitudine quam ostendit tabula.



84



495

THE  
SLENDER-LEGGED CRAB.



*GENERIC CHARACTER.*

*Legs* generally eight (in some species six or ten,) besides two claspers or chelated arms.

*Feelers* six, unequal.

*Eyes* two, generally distant, footstalked, moveable.

*Tail* articulated.

*SPECIFIC CHARACTER, &c.*


Small CRAB with heart-shaped tuberculated body, bifid snout, and very long slender hairy legs.

SLENDER-LEGGED CRAB.

*Penn. Brit. Zool.*

---

This curious species is distinguished from all others by the peculiar length and slenderness of its limbs, which give it at first view more the appearance of a Spider or a Phalangium than of a Crab. It is a native of the European coasts, and is generally of the size represented on the plate.



## PICUS PRINCIPALIS.



### CHARACTER GENERICUS.

*Rostrum* polyedrum, rectum, apice cuneato.

*Nares* pennis setaceis recumbentibus obtectæ.

*Lingua* teres, lumbriciformis, longissima, mucronata, apice retrorsum aculeata setis.

*Pedes* scansorii.

*Lin. Syst. Nat.* p. 173.

### CHARACTER SPECIFICUS, &c.

PICUS niger, crista coccinea, linea utrinque col-  
lari remigibusque secundariis albis.

*Lin. Syst. Nat.* p. 173.

PICUS niger Carolinensis.

*Briss. av.* 4. p. 26.

PICUS imbrifoetus.

*Raii syn.* p. 162.

---

Inter Picum principalem seu albirostem et illum qui pileatus dicitur summa videtur esse affinitas. Generantur ambo in iisdem Americæ septentrionalis partibus, et eadem fere sunt indole. Plurimis harum avium rostris orbiculatim dispositis feruntur

B . . . . . principes



principes Americani corollarum seu diadematum  
margines ornare. Superat pileatum magnitudine  
Picus principalis. Depingitur in tabula quasi tertia  
pars veræ mensuræ.





R.N.

London Published Oct 17 1801 by Elizabeth Nodder Juno & Co.

Newman Street

THE  
WHITE-BILLED WOODPECKER.



*Bill* polyedral, strait, wedge-shaped towards the tip.

*Nostrils* covered by recumbent setaceous feathers.

*Tongue* worm-shaped, very long, sharp-pointed, and edged towards the tip with reversed bristles.

*Feet* scansorial.

*SPECIFIC CHARACTER, &c.*

Black WOODPECKER with scarlet crest and white neck-stripe and secondaries.

WHITE-BILLED WOODPECKER.

*Catesb. Carol.* 1. pl. 16.

Pic à bec blanc.

*Buff.* 7. p. 46. Pl. Enl. 690.

---

The *Picus principalis* or White-billed Woodpecker, extremely nearly allied to the *Picus pileatus*, is a native of the same parts of North America, and is of similar habits. It is said that the Indian chiefs

hold the bill of this bird in high estimation, ornamenting their crowns or diadems with several of the beaks set in a circular manner round the margin. In size it is considerably larger than the pileatus: the figure represents it reduced about two thirds from its natural dimensions.

## SCOMBER DUCTOR.



### CHARACTER GENERICUS.

*Caput* compressum, læve.

*Membrana branchiostega* radiis septem.

*Corpus* læve, linea laterali postice carinatum.

*Pinnæ spuriae* sæpius versus caudam.

*Lin. Syst. Nat.* p. 492.

### CHARACTER SPECIFICUS, &c.

SCOMBER glaucus, fasciis transversis cyaneis,  
cauda fasciis duabus obliquis nigris.

Gasterosteus DUCTOR. G. spinis dorsalibus  
quatuor.

*Lin. Syst. Nat.* p. 489.

## SCOMBER DUCTOR.

*Bloch. ichth.* 10. p. 51. t. 338.


---

Quod squalo iter facienti comitem se sæpissime  
adjungere et præire solitus sit, non inepte inditum  
est huic pisci nomen *Sombri Ductoris*. Incolit præ-  
cipue mare Mediterraneum et Atlanticum, Sombro  
communi paulo major, et in cibis habetur delica-  
tioribus.









London, Published Oct 1830, by Elizabeth Nodden, Low of Newman Street.

B.V.

THE  
PILOT MACKREL.



*GENERIC CHARACTER.*

*Head* compressed, smooth.

*Gill-Membrane* furnished with seven rays.

*Body* smooth, carinated at the hind part by the lateral line.

*Spurious Pinnules* (in many species) towards the tail.

*SPECIFIC CHARACTER, &c.*

Glaucous MACKREL, with deep-blue transverse bands, and two oblique black bands across the tail.

The PILOT-FISH.

The PILOT MACKREL.

---

The Scomber Ductor or Pilot Mackrel, so named from its often preceding or accompanying the Shark in its excursions, is principally seen in the Mediterranean and Atlantic seas. In size it is somewhat superior to a common Mackrel, and is considered as an excellent article of food.



## STROMBUS FUSUS.



### CHARACTER GENERICUS.

*Animal* Limax.

*Testa* univalvis, spiralis, latere ampliata.

*Apertura*, Labro sæpius dilatato, desinens in  
canalem sinistrum.

*Lin. Syst. Nat.* p. 1207.

### CHARACTER SPECIFICUS, &c.

STROMBUS testa turrita lævi, cauda subulata,  
labro dentato.

*Lin. Syst. Nat.* p. 1207.

Murex FUSUS.


*Lin. Mus. Lud. Ulr.* 638. No. 316.

*Seb. mus.* 3. t. 56. f. 1. 2. 3.

---

In littoribus Americanis præcipue conspicitur  
Strombus Fusus, inter conchylia rariora numerandus:  
magnitudinem generalem ostendit tabula.







R.N. London, Published Sep. 7<sup>th</sup> 1800 by Elizabeth Nodder, Sons & Co. Newman Street.

THE  
SPINDLE STROMBUS.



*GENERIC CHARACTER.*

*Animal* allied to a *Limax*.

*Shell* univalve, spiral, amplified on one side.

*Aperture* with the lip generally dilated, ending  
in a channel towards the left.

*SPECIFIC CHARACTER, &c.*

Yellowish-Brown **STROMBUS**, with smooth  
turrated shell, subulated tail, and toothed  
lip.

Fuseau étoilé épais.

*Knorr. vergn. 5. pl. 6. 7.*

Le Grand Fuseau blanc.

*Argenv. pl. 13. D.*

---

The Spindle Strombus is principally found about the American shores, and is numbered among the rarer shells: its general size is shewn on the annexed plate.





## PAPILIO AGENOR.



### CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatae.

*Alæ* (sedentis) erectæ sursumque conniventes, (volatu diurno.)

*Lin. Syst. Nat.* p. 744.

### CHARACTER SPECIFICUS, &c.

PAPILIO alis dentatis nigris basi sanguineis, primoribus striatis, posticis disco albo maculis nigris.

*Lin. Syst. Nat.* p. 747.

*Clerk. ic. ins. rar.* t. 15.

*Cram. pap.* 3. t. 32. f. A. B.

*Jablonsky pap.* 1. t. 8. f. 3.

---

In variis Asiæ regionibus generatur Papilio Agenor, magnitudine vera in tabula depictus.







London, Published Sept. 17<sup>th</sup> 1801, by Elizabeth, Mother, to the Rev. Thomas Martineau Street.

## AGENOR.



### *GENERIC CHARACTER.*

*Antennæ* commonly thickening towards the end into a clavated tip.

*Wings* (when at rest) meeting upwards. (Flight diurnal.)

### *SPECIFIC CHARACTER.*

Black Butterfly with dentated wings red at the base; the lower pair white on the disk, with black spots.

---

This elegant insect is found in several of the Asiatic regions, and is represented on the plate in its natural size.



## LEMUR CALAGO.



### *CHARACTER GENERICUS.*

*Dentes Primores* superiores quatuor : intermediis remotis. Inferiores sex : longiores, porrecti, compressi, paralleli, approximati.

*Laniarii* solitarii, approximati.

*Molares* plures, sublobati : antichi longiores, acutiores.

### *CHARACTER SPECIFICUS, &c.*

LEMUR CALAGO. L. caudatus albidus, subtus griseus, cauda ferruginea.

CALAGO.

*Geoffroy Magaz. Encycl.* 1. p. 41. pl. 1.

---

Senegaliæ incolit rarum hoc animal quod primum detexisse censeo Adansonum. In arboribus versatur, quarum in cavis prolem alere dicitur, nido foliis et gramine intus stipato. Insectis præcipue vescitur quæ manibus perite arrepta celerrime devorat. Magnitudo ei est quasi sciuri vulgaris. Color ex albo flavet, tinctura quadam cinerea subtus adumbratus. Cauda ferruginea, caput canum. Aures prægrandes, tenues, erectæ, apicibus rotundatis, in-



tus roseæ. Ingenio esse dicitur miti et innocuo, et in cibus esse Galam et alias regiones Africanas incolentibus. Inter hunc et illum quadrupedem quem nomine *Fennec* descripsit Dominus Brucius, evidens omnino et manifesta est similitudo.





London. Published & Sold at 1801 by Elizabeth. Author, John G. Vancouver Street.

R. V.

THE  
CALAGO LEMUR.



*GENERIC CHARACTER.*

*Front-Teeth* in the upper jaw four: the intermediate ones remote.

In the lower jaw six: longer, stretched forwards, compressed, parallel, approximated.

*Canine-Teeth* solitary, approximated.

*Grinders* several, sublobated: the foremost somewhat longer and sharper.

*SPECIFIC CHARACTER, &c.*

Long-tailed whitish LEMUR, grey beneath, with ferruginous tail.

*Gen. Zool.* 1. p. 108.

CALAGO.

*Audeb. Singes, &c.*

---

This rare animal is a native of Senegal, and seems to have been first discovered by Adanson. It inhabits trees, in the hollows of which it is said to prepare its nest, lining it with leaves and grass. It feeds principally on insects, which it catches with wonderful dexterity. Its size is nearly that of a

C 2


common.

common squirrel, and its colour yellowish white, with a slight cast of ash-colour on the under parts: the head is grey, and the tail ferruginous: the ears very large, thin, upright, rounded at their extremities, and internally of a bright rose-colour. It is said to be of a mild and gentle disposition, and is used as an article of food by the natives of Galam and some other parts of Africa. There is a remarkable similarity in point of general appearance between this animal and the *Fennec*, described by Mr. Bruce.



## CONUS ARACHNOIDEUS.

122



*CHARACTER SPECIFICUS, &c.*

CONUS testa spadiceo reticulata, fasciis duabus  
tribusve obscurioribus, spira coronata acuta.

*Lin. Syst. Nat. Gmel. p. 3388.*


*Knorr. Vergn. 6. t. 4. f. 4.*

*Spengl. selt. conch. t. 1. f. D.*

*Martini conch. 2. t. 61. f. 676.*

---

In oceano Indico generantur conchyliā pulcher-  
rīma magnitudine naturali in tabula depicta.







R.N

London Published Novr 1<sup>o</sup> 1801, by Elizabeth Nodder, Sons & Newman Street.

THE  
ORANGE-FLAG CONE  
AND  
COBWEB CONE.



*GENERIC CHARACTER.*

*Animal* resembling a Limax or Slug.

*Shell* more or less conic or pyramidal.

*Aperture* longitudinal, linear, without teeth,  
entire at the base.

*Pillar* smooth.



ORANGE-FLAG CONE.



*SPECIFIC CHARACTER, &c.*

CONE with smooth flesh-coloured shell, with  
whitish bands, and the upper spires chan-  
nelled.

The speckled ORANGE-FLAG CONE.

COBWEB

## COBWEB CONE.



### *SPECIFIC CHARACTER, &c.*

White CONE with chesnut-coloured reticulations, two or three dusky bands, and sharp, coronated spire.

The SPIDER CONE, or COBWEB CONE.

---

The beautiful shells represented in their natural size on the present plate are natives of the Indian seas.

# ANABLEPS TETROPTHALMUS.



## CHARACTER GENERICUS.

*Corpus* cylindricum.

*Os* dentatum.

*Bloch. ichth.* 11. p. 3.

*Abdominales.*

## CHARACTER SPECIFICUS, &c.

ANABLEPS oculis prominentibus, pupillis duplicatis.

ANABLEPS oculis prominentibus, cirris duobus.

*Bloch. ichth.* 11. p. 5.

Cobitis ANABLEPS. C. cirris duobus, capite depresso, oculis prominulis.

*Lin. Syst. Nat.* p. 499.

ANABLEPS.


*Artedi gen. pisc.* 25.

---

Mirum aliquid et insolitum contigit huic pisci, quod nempe in una eademque orbita oculi sint quasi geminati. Ab accurata tamen dissectione satis probatum est oriri hoc e cornea tunica in duas partes  
divisa,

divisa, adeo ut pupilla utriusque oculi videatur veluti duplicata. Rem insignem dilucide explicat celeberrimus Blochius. Generatur Anableps tetropthalmus in maribus Americanis, et crescere solet in longitudinem pedalem.





London Published Nov. 17 1801 by Elizabeth Nodder Son of P. Newman Street.

THE  
FOUR-EYED ANABLEPS.



GENERIC CHARACTER.

- Body* cylindrical.
- Mouth* dentated.
- Eyes* prominent.

SPECIFIC CHARACTER, &c.

ANABLEPS with prominent eyes and double pupils.

Das Vierauge.

*Bloch ichth. t. 361.*

---

The fish represented on the present plate is distinguished by a highly remarkable particularity in its conformation ; the eyes appearing as if double. This however, on accurate dissection, is found to be owing to an internal division of the cornea, in such a manner as to present the appearance of two pupils in each eye ; a circumstance well explained in the work of Dr. Bloch. The *A. tetropthalmus* is a native of the American seas, and measures about twelve inches in length.





## PAPILIO PROTESILAUS.



### CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatae.

*Alæ* (sedentis) erectæ sursumque conniventes, (volatu diurno.)

*Lin. Syst. Nat.* p. 744.

### CHARACTER SPECIFICUS, &c.

PAPILIO alis caudatis subconcoloribus albidis, fasciis fuscis, unica subtus sanguinea, angulo ani rubro.

*Lin. Syst. Nat.* p. 752.

*Merian Surin.* t. 43.

*Seb.* 4. t. 36. f. 11. 12.

*Aubent. miscell.* t. 44. f. 1. 2.



*Cram. pap.* 17. t. 202. f. A. B.

---

Varias Americæ meridionalis regiones incolit Papilio Protesilaus, cujus magnitudinem veram ostendit tabula.







R. N.

London Published Nov 11. 4301. by Elizabeth Nodden Sons & Co. Newman Street.

## PROTESILAUS.



### *GENERIC CHARACTER.*

*Antennæ* commonly thickening towards the end  
into a clavated tip.

*Wings* (when at rest) meeting upwards. (Flight  
diurnal.)

### *SPECIFIC CHARACTER, &c.*

White Butterfly with tailed wings banded with  
brown, and with a crimson stripe beneath.

Page de la Reine.

*Merian Surin.* 43.

Le Flambé du Perou.

*Pl. Enl.* 44. f. 1. 2.

---

This most elegant insect is a native of several  
parts of South America, and is represented on the  
plate in its natural size.



## FRINGILLA ELEGANS.



### CHARACTER GENERICUS.

*Rostrum* conicum, rectum, acuminatum.

*Lin. Syst. Nat.* p. 317.

### CHARACTER SPECIFICUS, &c.

FRINGILLA olivacea, fronte gula caudaque rubris, abdomine albo nigroque undulato.

FRINGILLA supra viridis, collo cinereo, pectore flavo, capistro gula uropygio caudaque rubris, abdomine maculis lunatis albis.

*Lath. ind. orn.* p. 441.

### FRINGILLA ELEGANS.

*Lin. Syst. Nat. Gmel.* p. 912.

Le beau Marquet.


*Buff. ois.* 3. p. 497.

*Pl. Enl.* 203. f. 1.

---

Africam incolit pulcherrima hæc avis, cujus magnitudinem veram ostendit tabula.









THE  
VARIEGATED FINCH.

505



*GENERIC CHARACTER.*

*Bill* conic, strait, sharp-pointed.

*SPECIFIC CHARACTER, &c.*

Olive-green FINCH, with red front, throat and tail, and abdomen variegated with black and white.

Elegant FINCH.

*Lath. syn.* 3. p. 266.

---

This highly beautiful species is a native of Africa, and is represented on the annexed plate in its natural size.



# CORYPHÆNA CÆRULEA.



## CHARACTER GENERICUS.

*Caput* maxime truncato-declive.  
*Membrana branchiostega* radiis quinque.  
*Pinna* dorsalis longitudine dorsi.

## CHARACTER SPECIFICUS, &c.

CORYPHÆNA CÆRULEA, iridibus aureis.

CORYPHÆNA tota CÆRULEA.

*Bloch ichth.* 6. p. 120. t. 176.

CORYPHÆNA CÆRULEA.

*Lin. Syst. Nat.* Gmel. p. 1191.

---

Maria incolit Americana Coryphæna cærulea, cujus iconem archetypam a celeberrimo Plumiero centum circiter abhinc annos delineatam publicavit Blochius, qui eam in splendidissimum opus ichthyologicum imitando transtulit. In bipedalem crescere solet longitudinem.

1000

1000

1000

1000

1000

1000

1000

1000

1000

1000

1000

1000

1000

1000

1000

1000


1000

1000

1000


1000

1000





516



RN

THE  
BLUE CORYPHÆNA.

506



*GENERIC CHARACTER.*

*Head* sloping suddenly downwards.

*Gill-membranes* five-rayed.

*Dorsal Fin* the length of the back.

*SPECIFIC CHARACTER, &c.*

BLUE CORYPHÆNA with gold-coloured  
irides.

*Novacula cærulea.* The Blue-Fish.

*Catesb. Car. 2. t. 18.*

*Le Rasoir bleu.*

*Bloch. ichth. 6. t. 176.*

---

The *Coryphæna cærulea* is a native of the American seas, and grows to about the length of two feet. An elegant figure of this fish occurs amongst the drawings of the celebrated Father Plumier, by whom it was described in the preceding century: of this figure Dr. Bloch has availed himself in his splendid ichthyological publication.



# APLYSIA DEPILANS.



## CHARACTER GENERICUS.

*Corpus* repens, obvelatum membranis reflexis,  
clypeo dorsali pulmones obtegente.

*Foramen* laterale dextrum.

*Anus* supra extremitatem dorsi.

*Tentacula* quatuor, anterieus sita.

## CHARACTER SPECIFICUS, &c.

APLYSIA fusco-flavescens, clypeo dorsali sub-  
purpurascete.

## APLYSIA DEPILANS.

*Lin. Syst. Nat.* p. 1082.

Lernæa.

*Bohadsch de anim. marin.* p. 1. t. 1. 2. 3. 4.

Monstrum marinum singulare.

*Seb.* 3. t. 1. f. 8. 9.



---

Ob summam affinitatem possint fere cum limacibus  
conjugi quæ in hoc genere continentur animalia,  
et quasi limaces marini reputari. Species præcipua  
quam depinximus, cui color communis pallet livide  
fuscus, magnitudine multum variat, longa interdum  
sex vel octo uncias, plerumque autem multo minor.

Male

**Male olet, diciturque contrectata fluidum quiddam  
acre emittere, quod si in cutem inciderit, ruborem  
excitat: quod tamen non Aplysiæ solius est, com-  
mune enim est huic cum multis in Molluscorum ge-  
nere. Circa littora maris Mediterranei plerumque  
invenitur Aplysia depilans.**





RN

THE  
MEDITERRANEAN APLYSIA.



*GENERIC CHARACTER.*

*Body* repent, covered by reflected membranes  
and a dorsal shield.

A *Foramen* on the right side.

*Vent* situated at the extremity of the back.

Four *Tentacula* situated in front.

*SPECIFIC CHARACTER, &c.*

Yellowish-Brown APLYSIA, with a purplish  
cast on the dorsal shield.

Depilatory APLYSIA.

*Brit. Zool.* 4. p. 35. pl. 6.

The Sea Hare.

---

The genus *Aplysia* is so nearly allied to that of *Limax*, that it might almost be conjoined with it, and may be considered as a kind of marine Slug. The principal species, which is here represented, varies much in size, and is sometimes found of the length of six or eight inches: it is however more commonly found of a smaller size. Its general colour is a pale  
livid



livid brown. It is an animal of a disagreeable smell, and, when handled, is said to discharge a fluid of an acrimonious nature, which causes a degree of redness and inflammation on the skin: this however is a particular which is by no means confined to the genus *Aplysia*, but takes place in a still greater degree in several other animals belonging to the tribe *Molusca*. The *Aplysia depilans* is chiefly found about the coasts of the Mediterranean.

## VOLUTA ÆTHIOPICA.



528

### CHARACTER GENERICUS.

*Animal* Limax.

*Testa* unilocularis, spiralis.

*Apertura* ecandata, subeffusa.

*Columella* plicata.

### CHARACTER SPECIFICUS, &c.

VOLUTA testa emarginata ventricosa, spira coronata spinis fornicatis, apice papillari, columella quadriplicata.

*Lin. Syst. Nat.* p. 1195.

Cymbium mamillare coronatum.

*Seb.* 3. t. 65. f. 12.


Corona ÆTHIOPICA.


*Argenv.* t. 20. f. F.

---

Cochlea hæc magna et venusta, quæ interdum grandior longe est quam in tabula ostenditur, e maribus Indicis extrahitur, colore, ut plurimum, fusco-flavescente; interdum, (quod et specimini huic nostro contigit) non sine fascia seu fasciis macularum magnarum albarum a forma cordis non magnopere distantium, aut angulatarum. Interdum quoque spinarum basin cingentium variat longitudo.







R.V.

THE  
ÆTHIOPIAN VOLUTE.



*GENERIC CHARACTER.*

*Animal* resembling a Limax or Slug.

*Shell* unilocular, spiral.

*Pillar* twisted or plaited.

*SPECIFIC CHARACTER, &c.*

Yellow-Brown, emarginated, ventricose VOLUTE, with four wreaths on the pillar, and the spire crowned with spines.

La Couronne d'Ethiopie.

*Argenv.* 17. f. F.

The ÆTHIOPIAN Crown.

---

This large and elegant shell, which sometimes far exceeds the size of the annexed engraving, is a native of the Indian seas. Its general colour is a yellowish brown, which is sometimes varied, as in the present specimen, with one or more transverse fasciæ of large and somewhat heart-shaped or angular spots: it also occasionally varies in having either longer or shorter spines or processes round the base.



## CORACIAS SINENSIS.



### CHARACTER GENERICUS.

*Rostrum* cultratum apice incurvato, basi pennis denudatum.

*Lingua* cartilaginea, bifida.

*Pedes* ambulatorii.

*Lin. Syst. Nat.* p. 159.

### CHARACTER SPECIFICUS, &c.

CORACIAS viridis, subtus albo-flavescens, fascia oculari alisque nigricantibus, cauda cuneiformi apice alba.

CORACIAS viridis, subtus viridi-flavicans, fascia per oculos alisque nigricantibus, remigibus secundariis caudaque cuneiformi apice albis.

*Lath. ind. orn.* p. 171.

CORACIAS viridis, subtus ex flavicante alba, cauda cuneiformi apice alba.

*Lin. Syst. Nat.* Gmel. p. 381.

Galgulus SINENSIS.

*Briss. av. 2.* p. 77. t. 6. f. 2.


---


Sinam incolit elegantissima hæc avis, a celeberrimo Brissonio primum descripta: magnitudo ejus est quasi Corvi glandarii.

E









RV

304

THE  
CHINESE ROLLER.



*GENERIC CHARACTER.*

*Bill* strait, bending a little towards the tip ;  
cultrated at the edges.

*Nostrils* narrow, naked.

*Legs* generally short : toes divided to their ori-  
gin ; three forwards and one backwards.

*SPECIFIC CHARACTER, &c.*

Green ROLLER, yellowish white beneath, with  
blackish wings and eye-stripe, and cuneated  
tail with white tip.

CHINESE ROLLER.

*Lath. syn.* 1. p. 414.

Rolle de la Chine.

*Buff. ois.* 3. p. 132.

Rollier de la Chine.

*Pl. Enl.* 620.

---

This highly elegant bird is a native of China, and  
seems to have been first described by the celebrated  
Brisson ; its size is that of a common Jay.



## TETHYS FIMBRIA.



### CHARACTER GENERICUS.

*Corpus* liberum, oblongiusculum, carnosum, apodum.

*Os* proboscide terminali cylindrica sub labio explicato.

*Foramina* duo ad latus colli sinistrum.

*Lin. Syst. Nat.* p. 1089.

### CHARACTER SPECIFICUS, &c.

TETHYS labro crenulato.

*Lin. Syst. Nat.* p. 1089.

FIMBRIA.

*Bohadsch de an. marin.* p. 54. t. 5.

---

Cum luculente adeo descriperit hoc animal celeberrimus Bohadsch in opere cui titulus *De Animalibus marinis* ut nulla opus sit emendatione, liceat mihi ipsius auctoris verba subjungere.

“ Candore, excepto labii margine, undique nitet, et longitudine sex pollices assequitur. Labium in anteriore capitis parte membranæ fimbriatæ adinstar expansum non exiguo animalis ornamento est, quod latitudine quatuor pollices cum dimidio adæquat, longitudine,

gitudine vero tres pollices non excedit. Margo ejus utrinque crenatus est, et crassior reliqua labii portione, supra quam utrinque eminent, non secus ac fimbria aurea vel argentea pileo circumdata. Unde etiam aliud quid quam continuatio membranæ reliquam labii portionem constituentis esse videtur. Color marginis fimbriati ex nigro et luteo variegatus est, ita ut crenata ejus interior portio nigra sit, punctis nonnullis luteis notata, huic vero opposita et quoque crenata portio ex integro nigricans; pars intermedia aureo veluti colore splendet. Elegans hæc colorum varietas in eo duntaxat marginis latere observatur, quod prono animalis situ correspondet, in opposito enim latere totus margo nigro depictus est colore. Membrana reliquum labii corpus constituens ex fibris crassis candidis fere tendineis compacta est. Ad initium capitis ubi nimirum labium mox descriptum suam ducit originem, duo tentacula auriformia plana, lata, nullo sinu aut cavitate prædita, quatuor lineas larga et sex lineas longa exurgunt, quæ figura sua aures canis investigatorii non inepte referunt. Oculi nulli ad radicem horum tentaculorum, neque etiam in ulla alia capitis parte visibiles occurrunt; qui si adessent eos in corpore candidissimo saltem lente detegere impossibile non fuisset. Retro tentacula auriformia dorsum initium suum sumit; quod sensim gracilescens pyramidem seu conum præ se fert; longitudo ejus tres pollices, et sex lineas comprehendit, et basis unum pollicem duasque lineas in diametro adæquat. Latera dorsi duplici ordine appendicum carnearum candidarumque quæ partim conicam partim cylindricam formam



formam habent, exornata sunt. Maximæ ex his appendicibus quinque, minimæ duas lineas longæ sunt. Præter has variæ protuberantiæ in dorso observantur, ex quibus communiter una alterave appendix propullulat. Ad utrumque dorsi latus alia tubera majora collocantur, quæ abdominis, dorso multum amplioris, portiones sunt.”

Circa oras maris Mediterranei præcipue conspicitur Tethys Fimbria.






No



519



ferent nature from the rest of the lip: the colour of this fimbriated part or edge is variegated with black and yellow, which variegations are most conspicuous on the concave or under part: the membrane constituting the expanded part of the lip is composed of strong white and almost tendinous fibres: at the beginning of the head, where the lip commences, are two ear-shaped tentacula, broad, flat, without any cavity, about six lines in length and four in breadth; and bearing some resemblance to the ears of a hound: no appearance of eyes can be distinguished, even by the assistance of a glass: the back, which commences immediately beyond the tentacula, is of a pyramidal figure or gradually lessening to a point: its length is about three inches and a half, and its breadth at the base something more than one inch: on each side the back down its whole length runs a series of white, fleshy, conical protuberances, the largest of which measure about five and the smallest two lines in length: besides these are also some scattered tubercles or appendices on the middle of the back; while on each side are certain swellings larger than the rest, and which are caused by the projecting parts of the abdomen which is of much greater diameter than the back.

This animal is principally found about the coasts of the Mediterranean.

## ECHENEIS REMORA.



### *CHARACTER GENERICUS.*

Caput supra planum, marginatum, transverse sulcato-serratum.

### *CHARACTER SPECIFICUS, &c.*

ECHENEIS cauda bifurca, striis capitis octodecim.

*Lin. Syst. Nat. p. 447.*  
*Thoracici.*

ECHENEIS cauda bifurca.

*Gronov. zooph. p. 75.*

REMORA.

*Rondel. Willughb. &c.*

---

Insignit hoc genus capitis area quasi plana et ovata, dissepimentis plurimis transversim divisa, qua a parte adjungere se solet Echeneis imo aut lateribus navium, nec non ipsis etiam piscibus majoribus. Auxit hoc in miraculum veterum superstitio, creditumque olim est posse illam navim pleno cursu volantem de subito impedire. Huic absurdissimæ opinioni ansam dedisse facile crediderim parvulam quandam primævarum gentium scapham plurimis hisce piscibus

piscibus simul adhærentibus paulisper remoratam et in latus deflexam ; et rumorem in vulgus sparsum a veris initiis crevisse tandem, ut fieri plerumque solet, in ridiculas fabulas ; quasi ingenita esset ipsi pisci naturalis quædam vis quæ navigiis ad libitum moræ posset esse et impedimento.

Innascitur Echeneis Remora in mari Mediterraneo et Atlantico, longa, ut plurimum, quindecim uncias.

THE  
FORK-TAILED REMORA.



*GENERIC CHARACTER.*

*Head* flattened at the top into an oval shield marked by numerous transverse divisions.

*SPECIFIC CHARACTER, &c.*

REMORA with forked tail and about eighteen divisions on the shield.


The REMORA or Pilot-Fish,

The Sucking-Fish.

---

This highly singular genus is at once distinguished by the uncommon appearance of the head, which is formed on the upper part into a flat, oval space, divided by numerous transverse dissepiments: by this part the animal adheres at pleasure either to the bottom or sides of vessels, or even to several of the larger fishes themselves. The popular superstition of the ancients magnified this into a kind of miraculous power in the animal, which was supposed to be able to stop a ship in full sail by adhering to it. This idea, absurd as it is, might yet have originated in truth; nor can it be thought improbable that  
some

511



R.N.





some small canoe, in the earlier ages of mankind, might in some degree have been impeded in its progress, or made to incline unequally by several of these fishes adhering to one side ; and the tale, once related, might have gradually grown into the exaggerated power afterwards ascribed to the animal. The usual length of this fish is about fifteen inches : it is a native of the Mediterranean and Atlantic seas.



PAP. PERANTHUS.

*Fabr. mant. ins. 2. p. 4 ?*

---

Ostenditur in tabula papilionum duorum perelegantium et exoticorum vera magnitudo, quorum qui altius depingitur Indiam, qui inferius, Cochinchinam incolit.

## ÆNEAS & PERANTHUS.



### GENERIC CHARACTER.

*Antennæ* commonly thickening towards the end into a clavated tip.

*Wings* (when at rest) meeting upwards. (Flight diurnal.)

### SPECIFIC CHARACTER, &c.

ÆNEAS. Black Butterfly with denticulated wings, the upper pair marked above by a green spot, the lower by a blood-red pal-mated spot.


PERANTHUS. Black Butterfly with denticu-lated tailed wings green at the base.

Green-shouldered Cochinchina Butterfly,

---

Of the two beautiful exotic Butterflies exhibited on the present plate the superior is a native of India, and the inferior of Cochinchina: both are represent-ed in their natural size.





## TROCHILUS PELLA.



### CHARACTER GENERICUS.

*Rostrum* subulato-filiforme apice tubulato, capite longius: Mandibula superior vaginans inferiorem.

*Lingua* filiformis, filis duobus coalitis tubulosa.

*Pedes* ambulatorii.

*Lin. Syst. Nat.* p. 189.

### CHARACTER SPECIFICUS, &c.

TROCHILUS curvirostris, rectricibus intermediis longissimis, corpore rubro, capite fusco, gula aurata, uropygio viridi.

*Lin. Syst. Nat.* p. 189.

*Polytmus surinamensis longicaudus ruber.*

*Briss. av. 3.* p. 690.

*Falcinellus gutture viridi.*


*Klein. av.* p. 108.


---


In maximis est sui generis splendida hæc avis, Americæ australis, præcipue Surinamiæ incola. Minor longe est feminæ nitor, nec eminentes habet sed æquales caudæ pennas duas intermedias, quæ mari reliquis multo longiores decus afferunt. Specimen eximium unde figura hæc nostra depicta est suppeditavit Museum Leverianum.













21

THE  
TOPAZ-THROATED HUMMING-BIRD.



*GENERIC CHARACTER.*

*Bill* filiform, tubular towards the tip, the upper mandible sheathing the lower.

*Tongue* filiform, consisting of two conjoined tubes.

*Feet* formed for walking.

*SPECIFIC CHARACTER, &c.*

Curve-billed rufous HUMMING-BIRD, with dusky head, topazine throat, and two very long intermediate tail-feathers.

Long-tailed red HUMMING-BIRD.


*Edw. pl. 32.*

TOPAZ HUMMING-BIRD.

*Lath. syn. 1. p. 746.*

---

This species, which is one of the largest of the Humming-Birds, is a native of the Southern parts of America, and, in particular, of Surinam. The female is much less brilliant than the male, and is distinguished by the want of the two long intermediate tail-feathers so conspicuous in the male. The beautiful specimen here figured is preserved in the Leverian Museum.



## HOLOCENTRUS TIGRINUS.



### CHARACTER GENERICUS.

Opercula squamata, serrata, aculeata.

*Bloch. ichth. 7. p. 45.*  
*Thoracici.*

### CHARACTER SPECIFICUS, &c.

HOLOCENTRUS maculatus, pinna caudæ lunata.

*Bloch. ichth. 7. p. 59. t. 237.*

Percæ maxilla inferiore longiore, toto maculis et lineis transversis varia.

*Seb. mus. 3. t. 27. f. 15.*


Prochilus capite producto, &c.

*Klein. misc. pisc. 5. t. 12.*

---

A Percæ genere, cui certe admodum affine est, præcipue dignoscitur Holocentri genus, quod operculorum spinosa sit facies. Speciem elegantem descripsimus, maria Indica incolentem, eadem ut plurimum magnitudine qua Clupea vulgaris sive *Harengus* Linnæi.





10

11



J.N.

THE  
VARIEGATED HOLOCENTRUS.



*GENERIC CHARACTER.*

Gill-Covers scaly, serrated, and spiny.

515,

*SPECIFIC CHARACTER, &c.*

White HOLOCENTRUS with brown variegations.

Tiger HOLOCENTRUS.

La Marquille.

*Renard Hist. Poiss. 1. t. 6. f. 45.*

---

The genus *Holocentrus* is extremely allied to that of *Perca*, differing only in the spiny appearance of the opercula or gill-covers. The elegant species represented on the plate is a native of the Indian seas, and is generally of the size of a common Herring.



# NAUTILUS POMPILIUS.



## CHARACTER GENERICUS.

*Animal* (Rumph. mus. t. 17. f. B.)

*Testa* univalvis isthmis perforatis concamerata,  
polythalamia.

*Lin. Syst. Nat.* p. 1161.

## CHARACTER SPECIFICUS, &c.

NAUTILUS testæ apertura cordata, anfractibus  
obtusis lævibus.

*Lin. Syst. Nat.* Gmel. p. 3369.

Cochlea margaritifera.

*Rondel. aq.* p. 97.

NAUTILUS major crassus.

*Rumph. mus.* t. 17. f. A. c.


---

Non possit non contemplantis animum percellere,  
quæ leviter interdum variata per universum genus  
pervadit Nautili Pompilii interna conformatio; cujus  
singula camera seu divisio cum illa quæ proxime  
adjacet, tubulo brevi et aperto connectitur, cellula  
prima et præcipua, cæteris multo ampliore, habita-  
culum, ut videtur, corpori animalis incolentis præ-  
bente;

bente; cujus animalis vera natura usque in hunc diem non satis accurate explorata est. Licet inditum sit huic testæ nomen *Nautili* genericum, non tamen in aquis navigare potest Pompilius, more istius quæ *Argonauta Argo*, seu *Nautilus papyraceus* communiter nominatur; cum qua tamen a nonnullis scriptoribus confundi videtur: Crescere non raro solet in magnam molem *Nautilus Pompilius*, in mari-bus Indicis præcipue repertus.







RN.

THE  
GREAT NAUTILUS.



*GENERIC CHARACTER.*

*Animal* not perfectly known.

*Shell* univalve, divided internally into a great many concentric cells, communicating by short tubes.

*SPECIFIC CHARACTER, &c.*

Whitish NAUTILUS with yellow-brown transverse bands, smooth spires, and cordated aperture.

The great pearly NAUTILUS.

The Chambered NAUTILUS.

---

The curious structure of this shell, (and which also runs, with some variation, thro' the whole genus,) cannot be contemplated without admiration; each cell or camera communicating with the next by a small and short open tube, the first or principal cell being far larger than the rest, and appearing destined to contain the chief part or body of the inhabiting animal, the nature of which is not yet properly understood. The Nautilus Pompilius is a  
native

native of the Indian seas, and frequently arrives at a very considerable size. Notwithstanding its title, the Nautilus does not possess the power of sailing on the surface in the manner of the Argonauta Argo, or Paper-Nautilus, with which it sometimes appears to be confounded in the works of authors.

## PAPILIO PANTHOUS.



### CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatae.

*Alæ* (sedentis) erectæ sursumque conniventes, (volatu diurno.)

*Lin. Syst. Nat.* p. 744.

### CHARACTER SPECIFICUS, &c.

PAPILIO alis dentatis nigris concoloribus, primoribus albo maculatis; posticis maculis albis nigra foetis.


*Lin. Syst. Nat.* p. 748.

PAPILIO Amboinensis, &c.

*Seb. mus.* 4. t. 44. f. 22.

---

Papilionem Panthoum in vivario hoc nostro antea descripsimus. Ostenditur in tabula magnitudine naturali quæ hujus habetur esse femina, ex splendidissimo Sebæ opere desumpta, quæque fortasse omnes papiliones quos adhuc novimus magnitudine superat.







## PANTHOUS.



### *GENERIC CHARACTER.*

*Antennæ* commonly thickening towards the end into a clavated tip.

*Wings* (when at rest) meeting upwards. (Flight diurnal.)

### *SPECIFIC CHARACTER, &c.*


**Dusky Butterfly**, with dentated wings spotted with white, the spots on the lower wings marked by a black patch.

**The Great dusky Amboina Butterfly.**

---

In this plate is exhibited, (from the work of Seba,) the supposed female of the *Papilio Panthous*, before represented in the present publication. The insect is shewn in its natural size, and may perhaps be considered as the largest of all the papilionaceous tribe.





## TROCHILUS SUPERBUS.



### CHARACTER GENERICUS.

*Rostrum* subulato-filiforme, apice tubulato, capite longius : *Mandibula* superior vaginans inferiorem.

*Lingua* filiformis, filis duobus coalitis tubulosa.

*Pedes* ambulatorii.

*Lin. Syst. Nat.* p. 189.

### CHARACTER SPECIFICUS.

**TROCHILUS** rectirostris viridi-aureus, vertice cæruleo, fascia per genas duplici nigro-alba, gula pectoreque phoeniceis.

---

De pulcherrima hac specie nihil pro certo cognitum est, nisi quod, ut fere solet reliquum genus, Americæ partes australes incolat. Ostendit tabula veram avis magnitudinem, a nemine, ut opinor, antea descriptæ aut depictæ.







THE  
STRIPE-CHEEKED HUMMING-BIRD.



*GENERIC CHARACTER.*

517

*Bill* filiform, tubular towards the tip, the upper mandible sheathing the lower.

*Tongue* filiform, consisting of two conjoined tubes.

*Feet* formed for walking.

*SPECIFIC CHARACTER.*

Strait-Billed gold-green HUMMING-BIRD,  
with blue-crown, double black-and-white  
check-stripe, and crimson throat and breast.

---

Of this most beautiful species, which appears to have been hitherto undescribed, nothing more is known than that, like most others of this genus, it is a native of the Southern parts of America. It is represented on the plate in its natural size.



## FISTULARIA TABACARIA.

\*\*\*\*\*

### CHARACTER GENERICUS.

*Rostrum* cylindricum, apice maxillosum.

*Membrana* branchiostega radiis septem,

*Corpus* elongatum.

### CHARACTER SPECIFICUS, &c.

FISTULARIA fusca cæruleo maculata, abdomine albido, pinnis rubris, cauda bifida setifera.

*Abd:inales.*

FISTULARIA TABACARIA. F. cauda bifida setifera.

*Lin. Syst. Nat. p. 515.*

Solenostomus, &c.

*Klein M. Pisc. 4. p. 25. n. 4.*

---


Insignit *Fistulariæ* genus forma rostri et longitudo corporis. In eo duæ tantum continentur species, quarum quæ depingitur fortasse vulgatio est, in mari Indico reperta, longitudine fere tripedali. Adjicitur tabulæ varietas hujus piscis, cui rostrum serratum, a celeberrimo Blochio descripta.







515



515





# STROMBUS LATISSIMUS.



## CHARACTER GENERICUS.

*Animal* Limax.

*Testa* univalvis, spiralis, latere ampliata.

*Apertura* labro sæpius dilatato, desinens in canalem sinistrum.

*Lin. Syst. Nat.* p. 1207.

## CHARACTER SPECIFICUS, &c.

STROMBUS labro rotundato maximo, ventre inermi, spira subnodosa.

*Lin. Syst. Nat.* p. 1211.

*Alata lata.*

*Rumph. t. 36. f. L.*

*List. conch. t. 856. Martini conch. t. 82. f. 832.*

*Alata lata.*

*Seb. 3. p. 163. t. 63. f. 1. 2.*

---

Labio seu latere oris exteriori latius expanso distinguitur Strombi genus, et spargitur in plurimas species. E rarioribus est illa quam selegimus, in maribus Indicis, ut plurimum, reperta, sæpius major longe specimine depicto.









RN

THE  
WING STROMBUS.



*GENERIC CHARACTER.*

519

*Animal* resembling a Limax.

*Shell* univalve, spiral, enlarged on the side.

*Aperture* generally with a dilated lip, ending in a channel towards the left.

*SPECIFIC CHARACTER, &c.*

**STROMBUS** with very large rounded lip, smooth body, and subnodose spire.

**Wide-lipped Indian STROMBUS.**

**WING STROMBUS.**

---

The genus Strombus is distinguished by the remarkable width or dilatation of the exterior lip or edge of the shell: the species are numerous, among which that represented on the present plate may be considered as one of the rarest: it is a native of the Indian seas, and is occasionally found of a much greater size than the specimen figured.



## PHALÆNA MACROURA.



### CHARACTER GENERICUS.

*Antennæ* setaceæ, a basi ad apicem sensim attenuatæ.

*Alæ* (sedentis) sæpius deflexæ (volatu nocturno.)  
*Lin. Syst. Nat.* p. 808.


### CHARACTER SPECIFICUS, &c.

PHALÆNA ferruginea, alis inferioribus caudatis longissimis.

*Drury ins.* 3. p. 39. t. 29.

---

Ex omnibus lepidopteris quæ hactenus physicis innotuerunt, nescio an non Phalæna macroura præcipue notatu digna sit ob enormem longitudinem appendicum, quasi in caudæ similitudinem excrescentium, quibus alæ inferiores terminantur. Africæ incola in phalænis rarioribus exoticis numeratur. In eleganti Domini Drurii opere censeo eam primo descriptam fuisse et depictam.







Rur

THE  
LONG-TAILED PHALÆNA.



GENERIC CHARACTER.

*Antennæ* setaceous, gradually lessening from the base to the tip.

*Wings* (when at rest) commonly deflected.  
(flight nocturnal.)

SPECIFIC CHARACTER, &c.

Ferruginous PHALÆNA, with the lower wings extended into extremely long tail-like processes.

*Drury ins.* p. 39. pl. 29.

---

Of all the lepidoptera yet known, the present species is perhaps the most remarkable for the excessive length of the tail-like processes or appendices with which the lower wings are terminated: it is a native of Africa, and is numbered among the rarest of the exotic Phalænæ. It seems to have been first described and figured in the elegant work of Mr. Drury.





## LANIUS BICOLOR.



### CHARACTER GENERICUS.

*Rostrum* rectiusculum, dente utrinque versus  
apicem, basi nudum.

*Lingua* lacera.

*Lin. Syst. Nat.* p. 134.

### CHARACTER SPECIFICUS, &c.

LANIUS cæruleus, subtus albus, cauda sub-  
æquali.

LANIUS cauda subæquali, supra cæruleus sub-  
tus albus, capistro nigro.

*Lath. ind. orn.* p. 75.

Loxia Madagascarina.

*Lin. Syst. Nat.* 1. p. 306.

---


Species perpulchra hic depicta insulas Africanas  
præcipue autem Madagascariam incolit: femina a  
mare coloribus multo minus vividis distinguitur.











# HOLOCENTRUS SOGO.



## CHARACTER GENERICUS.

*Opercula squamata, serrata, aculeata.*

*Bloch. ichth. 7. p. 47.*

*Thoracici.*

## CHARACTER SPECIFICUS, &c.

HOLOCENTRUS ruber, lineis longitudinalibus  
flavis, cauda furcata.

HOLOCENTRUS SOGO. H. pinna ventrali  
radiis octo.

*Bloch. ichth. 7. p. 46. t. 232.*

HOLOCENTRUS.

*Seb. mus. 3. p. 73. t. 27. f. 1.*

Perca marina rubra.

*Catesb. Carol. 2. t. 3. f. 2,*


---

In maribus Indicis innascitur Holocentrus Sogo, magnitudine Percam vulgarem sive fluviatilem referens. Viro celeberrimo Carolo Plumier figuram archetypam hujus piscis debemus, a Blochio primum evulgatam.









RTN

THE  
RED HOLOCENTRUS.



*GENERIC CHARACTER.*

*Gill-Covers* scaly, serrated, spiny.

*SPECIFIC CHARACTER, &c.*

RED HOLOCENTRUS with longitudinal yellow stripes and forked tail.

The Sogo or RED HOLOCENTRUS.

The Squirrel.


*Catesb. Carol.* 2. pl. 3. f. 2.

Der Sogo.

*Bloch. ichth.* 7. pl. 232.

---

The Holocentrus Sogo is a native of the Indian seas, and is of the size of a common perch. It is to the celebrated Plumier that we owe the archetypal figure of this fish, which has been engraved in the work of Dr. Bloch.



## CONUS TULIPA.



### CHARACTER GENERICUS.

*Animal* Limax.

*Testa* univalvis, convoluta, turbinata.

*Apertura* effusa, longitudinalis, linearis, edentula, basi integra.

*Columella* lævis.

*Lin. Syst. Nat.* p. 1165.

### CHARACTER SPECIFICUS, &c.

CONUS testa ventricosa alba, nebulis longitudinalibus aurantiis, punctis transversis fuscis, apertura dehiscente.

CONUS TULIPA. C. testa oblonga gibba lævi, apertura dehiscente.

*Lin. Syst. Nat.* p. 1172.

*Rumph. mus.* t. 34. f. κ.

*Knorr vergn.* 5. t. 20. f. 1. 2.



---

Inter testas pulcherrimas merito numeratur Conus Tulipa cujus magnitudinem veram ostendit tabula. Circa littora Indica præcipue conspicitur.









R.D.Y.

THE  
TULIP CONE.



*GENERIC CHARACTER.*

*Animal* resembling a Limax or Slug.

*Shell* more or less conic or pyramidal.

*Aperture* longitudinal, linear, without teeth,  
entire at the base.

*Pillar* smooth.


*SPECIFIC CHARACTER, &c.*

Smooth white ventricose CONE, with longitudinal orange-coloured clouds, and dusky transverse specks.

The Tulip-Shell or TULIP CONE.

---

The Tulip Cone is justly considered as one of the most beautiful shells of its tribe: it is a native of the Indian seas, and is represented on the plate in its natural size.



## CANCER CRISTATUS?



### *CHARACTER GENERICUS.*

*Pedes* octo (rarius sex aut decem;) insuper manus duæ chelatae.

*Oculi* duo, distantes, plurimis pedunculati; elongati, mobiles.

*Cauda* articulata, inermis.

### *CHARACTER SPECIFICUS, &c.*

**Cancer** thorace pedibusque aculeatis, rostro porrecto bifido cristato.

*Fabr. ins.* 1. p. 503.

*Lin. Gmel.* p. 2980.

*Rumph. mus.* t. 8. f. 1.

*Seb.* 3. t. 22. f. 2. 3.


---


Hujus speciei descriptionem a Seba in opus nostrum transferre non dubitavimus.

“Miratu dignissimus iste cancer, Rumphio cancer spinosissimus dictus, non tantum in tegmine superiore, quod mire gibbosum, et quasi in montes et valles distinctum est, sed in aliis quoque partibus pilosa quasi tela, et pungentes aculeos, manipulatim compositos, ostentat. Caput, veluti turritum, acuminatum, spinosis processibus undique asperum est.

Brachia

Brachia itidem crassa, longa, ramos, in omni ambitu, ramorumque propagines emittunt. Quin vel ipsi forcipes similibus undique horrent. Octo postici pedes, pari modo constituti, aculeis crassis, pungentibus, simplicibus, ubique pleni sunt. Sic et cauda sese habet plurimis brevibusque articulis geniculata. Pedes posteriores in minutum et brevem terminantur unguiculum. Color dilute griseus per omnia regnat." Maria incolit Asiatica.





THE  
CRESTED CRAB.



GENERIC CHARACTER.

*Legs* generally eight, (in some species six or ten, besides two claspers or chelated arms.

*Eyes* two, commonly distant; footstalked, moveable.

*Tail* articulated.

SPECIFIC CHARACTER, &c.

Short-tailed tuberculated CRAB with numerous ramified and fasciculated spines, and bifid crested snout.

---

This highly singular species, says Seba, is the *cancer horridus* of Rumphius, and is covered not only on the upper part of the body, which is very convex and marked into various elevations and depressions, but also on every other part with groupes of sharp spines and hair-like prickles: the head is elevated, acuminate, and every where beset with spiny processes: the long and thick arms are covered along their whole outline with branching subdivisions which are visible even on the forcipes themselves: the eight hind legs are formed on a similar plan,



plan, but are covered with thick, sharp, simple spines: the tail, which is spiny also, consists of several small and short joints: the hind feet terminate in minute and short claws: the general colour of the whole animal is pale grey. It is a native of the Asiatic seas.

## ALCEDO CHLOROCEPHALA.



### *CHARACTER GENERICUS.*

*Rostrum* trigonum, crassum, rectum, longum.

*Lingua* carnosae, brevissima, plana, acuta.

*Pedes* gressorii plerisque.

*Lin. Syst. Nat.* p. 178.

### *CHARACTER SPECIFICUS, &c.*

ALCEDO viridi-cærulea, subtus alba, vertice viridi nigro marginato.

### ALCEDO CHLOROCEPHALA.

*Lath. ind. orn.* p. 250.

*Lin. Syst. Nat.* Gmel. p. 454.

---

Insulas Indicas inhabitat elegantissima hæc avis, magnitudine Alcedinem Europæam sive Ispidam paulum superans.







THE  
GREEN-HEADED KINGFISHER.



*GENERIC CHARACTER.*

*Bill* trigonal, thick, strait, long.

*Tongue* fleshy, very short, flat, sharp-pointed.

*Feet* gressorial.

*SPECIFIC CHARACTER, &c.*

Blue-Green KINGFISHER, white beneath, with  
green crown margined with black.

Le Martin-Pecheur a tete verte.

*Buff. ois.* 7. p. 190.

*Pl. Enl.* 783. f. 2.

GREEN-HEADED KINGFISHER.

*Lath. syn.* 1. p. 620.

---

This highly elegant species is a native of the Indian islands. Its size somewhat exceeds that of the common European Kingfisher.



## EQUES AMERICANUS.



### CHARACTER GENERICUS.

*Dentium* ordines plurimi.

*Corpus* fasciatum.

### CHARACTER SPECIFICUS, &c.

EQUES pinna dorsi altissima.

*Bloch. ichth.* 10. p. 76. t. 347.

*Thoracici.*

Chætodon lanceolatus. C. cauda integra, corpore fasciis tribus; oculari, pectorali, longitudinalique.

*Lin. Syst. Nat.* p. 466.

---

Piscem hunc in Chætodontibus a Linnæo habitum ad novum et sibi proprium genus retulit Blochius, quod dentium non sit, ut Chætodontum, unicus ordo, sed plures concentrici. Maria incolit Americana, et in tabula magnitudine quasi dimidiata depingitur.









189X

925

THE  
KNIGHT-FISH.



*GENERIC CHARACTER.*

*Teeth* in several rows.

*Body* banded.

*SPECIFIC CHARACTER, &c.*

KNIGHT-FISH with very high dorsal fin.

The Guaperva.

The Ribband-Fish.

*Edw. pl. 210.*

Le Chevalier.

*Bloch. ichth. t. 347.*

---

This fish, which by Linnæus was associated with the Chætodons, has been formed into a distinct genus by Dr. Bloch on account of its teeth, which, instead of being arranged in a single row, as in those animals, are disposed into several concentric ranges. It is a native of the American seas, and is represented on the plate of about half its natural size.



## MUREX RAMOSUS.



### CHARACTER GENERICUS.

*Animal* Limax.

*Testa* univalvis, spiralis, exasperata suturis membranaceis.

*Apertura* desinens in canalem integrum, rectum seu subadscendentem.

*Lin. Syst. Nat.* p. 1213.

### CHARACTER SPECIFICUS, &c.

MUREX testa trifariam frondosa, spira contigua, cauda truncata.

*Lin. Syst. Nat.* p. 1215.

## MUREX RAMOSUS.

*Rumph. mus.* t. 26.

*Purpura ramosa.*

*Argenv.* t. 16. f. c. E.

*List. conch.* t. 946. f. 41.

---


Contigit ferme omni Muricis generi ut obducatur testa exterior inæquali quadam et rugata scabritie: at insignis omnino est species quam describimus, quod spiræ seu volumina testæ processus efficiant plurimos

plurimos elongatos et dilatatos, apicibus magis minusve ramosis seu divaricatis.

Maria incolit Indica et Americana murex ramosus major interdum speciminibus quæ in tabula monstrantur.







THE  
BRANCHED MUREX.



*GENERIC CHARACTER.*

*Animal* allied to a Slug.

*Shell* univalve, spiral, roughened by membranaceous sutures.

*Aperture* ending in a strait or subascending channel.

*SPECIFIC CHARACTER, &c.*

Yellowish-brown transversely striated MUREX,  
with three longitudinal rows of foliaceous  
processes.

The BRANCHED MUREX.

Branched Purple-Shell.

---

The genus *Murex*, distinguished in general by a peculiar irregularity of surface, affords few species more remarkable than the present, in which the spires of the shell are beset with numerous lengthened and expanded processes more or less divided or branched at their extremities. It is a native of the Indian and American seas, and sometimes exceeds the size of the specimens represented on the annexed plate.



## SPHINX ATROPOS.



### CHARACTER GENERICUS.

*Antennæ* subprismaticæ, utroque fine attenuatæ.

*Lingua* exserta (plerisque.)

*Palpi* duo reflexi.

*Alæ* deflexæ.

### CHARACTER SPECIFICUS, &c.

SPHINX alis primoribus griseis, posterioribus abdomineque luteis nigro-fasciatis, thorace macula craniolari flava.

SPHINX alis integris: posticis luteis fasciis fuscis, abdomine luteo cingulis nigris.

*Lin. Syst. Nat.* p. 799.


---

Varias Europæ regiones incolit formosissimum hoc insectum, et e rarissimis est lepidopteris quæ in Britannia aluntur. Hujus larva solani tuberosi nec non jasmini folia præ cæteris amat, et cum noctu præcipue vescatur, interdiu deliteat, eo fit ut difficilior sit detectu quam aliæ pleræque larvæ lepidopterorum. Mense Augusto, chrysalidis vices subitura sub humo se condit, unde mense Septembri insequentis anni prodit ipsa Sphinx, thoracem habens macula

macula insignitum cuius possit esse quædam imaginaria similitudo cum cranio arido mortui hominis ut vulgo depingitur; quam ob causam, emissamque vocem stridulam, si quis tangendo irritaverit, indoctæ plebi magnum non raro timorem incutit quasi male ominata, et mortis prænuncia: immo narrat celeberrimus Reaumurius ipsas monachas Gallici cujusdam coenobii subitanea trepidatione fuisse percussas visa Sphinge quæ vespertina per fenestras involaverat.

Quod commune est aliis plerisque insectis, coloribus vel saturatoribus vel pallidioribus variat Sphinx Atropos.





RPN

THE  
JASMINE SPHINX.



*GENERIC CHARACTER.*

*Antennæ* subprismatic, attenuated at each extremity.

*Tongue* generally exerted.

*Feelers* two, reflex.

*Wings* deflected.

*SPECIFIC CHARACTER, &c.*

SPHINX with the upper wings grey, the lower wings and abdomen deep yellow barred with black, and the thorax marked by a scull-shaped yellow patch.

The Tiger Hawk-Moth.

The Jasmine Hawk-Moth, or Death-Head Sphinx.

---

This highly beautiful insect is a native of many parts of Europe, and is numbered among the rarer species of British lepidoptera. The larva or caterpillar is observed to prefer the leaves of the Jasmine and the Potatoe to those of most other plants; and as it feeds principally by night, concealing itself during the day, is much less frequently seen than most



most other insects of its tribe. It retires underground in the month of August, in order to undergo its change into a chrysalis, from which, in the month of September in the following year proceeds the complete insect, which is distinguished by a remarkable spot or patch on the thorax, bearing an imaginary resemblance to the figure of a scull or death's-head, as vulgarly represented. From this circumstance, joined to the stridulous tone which it utters when handled or irritated, it has been considered by the vulgar as an animal of ill omen, and as a messenger of fate. The celebrated Reaumur informs us that the members of a female convent in France were seized with general consternation on discovering one of these insects, which had accidentally flown in at one of the open windows during the evening. Like most other insects, it occasionally varies somewhat in colour, being darker or lighter in different individuals.

## PELECANUS SINENSIS.



### CHARACTER GENERICUS.

*Rostrum* rectum : apice adunco, unguiculato.

*Nares* rima oblitterata.

*Facies* nudiuscula.

*Pedes* æquibres: digitis omnibus quatuor simul palmatis.

### CHARACTER SPECIFICUS.

PELECANUS supra fuscus, subtus albidus fusco maculatus, gula alba, iridibus cæruleis, cauda rotundata.

---

In genere Pelecano nonnullæ sunt aves quæ ad exercendam artem piscatoriam dominis quæstuosam ali possint et institui. A Sinensibus deductus est mos, indeque reversi non ita pridem legati Britannici certiores fecere Europæos de specie qua utuntur ii ad hoc opus educata: illa nempe, quæ anserina quasi magnitudine, corpore tamen tenuior, si colores excipias, *graculo* et *carboni* pelecans non longe est absimilis.


Quando piscari volunt Sinenses, aves multas levibus in scaphis imponunt, quæ ad nutum domini cum in aquas se projecerint, parvo temporis spatio pisces


K

rostris

rostris arreptos in cymbam deferunt; nec annulo collari opus est ne forte prædam deglutiant (sine quo apud Europæos nunquam committitur *carboni* res piscatoria) adeo enim disciplinæ et imperiis assueverunt dociles Sinensium pelecani, ut si prædæ aliquid iis dispartitum fuerit, satis id putent ad laboris remunerationem.

Auctor est Dominus Stantonus, qui librum nuperime edidit de moribus Sinensium, conspici posse in nonnullis lacubus millia multa scapharum, unde mirum hoc piscaturæ genus ab avibus peragitur.





PLATE

THE  
CHINESE PELICAN.



*GENERIC CHARACTER.*

*Bill* long, strait, hooked at the end.  
*Nostrils* small, in an obscure furrow.  
*Face* naked.  
*Toes* all four united by a web.

*SPECIFIC CHARACTER.*

Brown PELICAN, with white throat, the body whitish beneath and spotted with brown; the tail rounded; the irides blue.  
*Staunton's Chinese Embassy*, 2. p. 388.

---

The art of training some particular species of this genus to the practice of fishing for the profit of their owners appears to have originated among that industrious people the Chinese, who are in the habit of turning every object to as much advantage as possible. It was not however till the period of the late embassy to China that the real species used for this purpose became known to Europeans. The bird is about the size of a common goose, but is of a thinner form, and resembles, except in colour, the European Shag and Corvorant. When used for fishing,

ing, these birds are carried in small boats, and, at a signal given, plunge into the water, and soon return with the prey in their mouth, requiring no ring round the neck, to prevent their swallowing it, as is the practice in Europe when the Cormorant is sometimes used for a similar purpose; the Chinese fowl being so well trained as to require only the encouragement of part of their prey. We are informed by Sir George Staunton, in his account of the late Embassy, that on some of the Chinese lakes may be seen thousands of small boats destined entirely to this curious species of fishery.

## PHALÆNA AGRIPPINA.



### CHARACTER GENERICUS.

*Antennæ* setaceæ, a basi ad apicem sensim attenuatæ.

*Alæ* (sedentis) sæpius deflexæ (volatu diurno.)

*Lin. Syst. Nat.* p. 808.

### CHARACTER SPECIFICUS, &c.

PHALÆNA (noctua) alis subconcoloribus albidis fusco nebulosis, subtus subviolaceis.

*Mer. ins. Sur.* t. 20.

## PHALÆNA AGRIPPINA.

*Cram. ins.* 8. t. 87. 88. A. A.

---


Phalænam hanc grandem et elegantem, Surinamiæ incolam, prima omnium descripsisse et depinxisse videtur celeberrima Merian in *historia insectorum Surinamensium*. Folia Gambogiæ depascitur larva, colore læte viridi, faciis latis, transversis, nigris variata: color autem viridis, anteaquam in chrysalidem convertatur phalæna, transire solet in splendido-rubrum. Errant frequenter auctores, quorum eadem est hæc species atque phalæna *strix* Linnæi, quæ




quæ tamen ad aliam tribum referri debet, generali quadam colorum dispositione huic affinis.


Jactat Museum Leverianum pulcherrima duo phalænæ Agrippinæ specimina.





RPN





RPN

## AGRIPPINA.



### *GENERIC CHARACTER.*

*Antennæ* setaceous, gradually diminishing from the base to the tip.

*Wings* (when at rest) commonly deflected.  
(Flight diurnal.)

### *SPECIFIC CHARACTER.*


Whitish Moth, clouded with brown variegations; the pattern nearly similar on both surfaces; the inferior tinged with violet-colour.

---

The large and elegant moth represented on the present plate is a native of Surinam, and appears to have been first described and figured by the celebrated Madam Merian, in her history of the insects of that country. The caterpillar or larva feeds on the leaves of the Gamboge-tree, and is of a fine green, variegated with large, transverse, black bands; but some time before its assuming the form of a chrysalis it is observed to exhibit a remarkable alteration of colour, the green changing to a beautiful red. This species is commonly, but mistakenly, quoted

quoted as the *Phalæna Strix* of Linnæus; that insect belonging to a different division of the genus, and being allied to the present species only in the general distribution of its colours. In the Leverian Museum may be seen two very fine specimens of the *Phalæna Agrippina*.

## ASCIDIA AURANTIUM.



### CHARACTER GENERICUS.

*Corpus* fixum, teretiusculum, vaginans.

*Aperturæ* binæ ad summitatem : altera humiliore.

*Lin. Syst. Nat.* p. 1087.

### CHARACTER SPECIFICUS, &c.

ASCIDIA subglobosa coccinea, papillis terminalibus cylindraceis.

ASCIDIA subglobosa, sacco coccineo punctis duriusculis scabro, papillis terminalibus cylindraceis rugosis.

*Pall. nov. act. Petrop.* 2. p. 246. t. 7.


*Lin. Syst. Nat. Gmel.* p. 3128.

Ab erudito domino Pallas mutuatus sum hanc descriptionem, qua nulla possit esse aptior et accuratior :

“ *Magnitudo* sæpe pomi aurantii majoris. *Forma*, præter basin truncatam testis lapillisque insidentem et papillas osculiferas, subglobosa. *Corium* externum in siccatis passim in magnas rugas crispatum, naturaliter æquabile, tenacissimum, rigidiusculum,



vix ungue crassius, extus totum punctis duriusculis, distantibus scabratum. *Papillæ* in vertice sphaeræ binæ cylindraceæ, rugosæ, altera major, utraque orificio cruciatim diffisso pervia. Intra cavum corii continetur *follis* ductibus duobus carnosus orificiis papillarum insertus, constans strato fibrarum extus circularium interioreque grossiorum longitudinalium, in discum baseos tendinosum, circularem convergentibus. Hic follis seu *ventriculus* facile integer a corio secedit et enucleatur, intus vacuus, aquam marinam recepturus, stipatus adnato *viscere* parenchymatoso, in anfractus intestiniformes efficto, flavescente, a basi per latus arcuato-adscendente. *Color* extus coccineus."





THE  
ORANGE ASCIDIA.



*GENERIC CHARACTER.*

*Body fixed, generally subcylindric.*

*Apertures two at the upper part; one lower than the other.*

*SPECIFIC CHARACTER.*

Subglobose orange-red ASCIDIA, with cylindrical terminal papillæ.

---

This species of *Ascidia* is described by Dr. Pallas, who informs us that it is a native of the Asiatic seas, and is often found of the size of a large orange. Its shape is nearly globular, but truncated at the base, where it is affixed to stones, shells, &c. the external skin or rind, which in the dried specimens is usually marked here and there by several large wrinkles, is, in its recent state, of an even surface, very tough, of scarcely more than a nail's thickness, and roughened all over with distant, callous specks: on the upper part are two cylindrical, wrinkled papillæ, one larger than the other, and both furnished with a crucial aperture: within the cavity of the rind is a sacculus or bladder, inserted by two fleshy ducts into the orifices of the papillæ, and consisting of circular exterior

exterior and longitudinal interior fibres, converging at the circular and tendinous base: this sacculus, which is for the reception of the sea-water, is easily separable from the rind: it is furnished internally with a yellowish visceral parenchyma, formed into intestinal windings ascending from the base up the sides of the cavity: the colour of this species is orange-red.

# UPUPA ERYTHORYNCHOS.



## CHARACTER GENERICUS.

*Rostrum* arcuatum, convexum, subcompressum,  
obtusiusculum.

*Lingua* obtusa, integerrima, triquetra, brevissima.

*Pedes* ambulatorii.

*Lin. Syst. Nat.* p. 183.

## CHARACTER SPECIFICUS.

UPUPA nigro-violacea, viridi nitens, rectricibus  
versus apices macula utrinque alba, rostro  
rubro.


*Cimelia Physica.* p. 96. t. 52.

UPUPA viridi-atra, purpureo splendens, &c.

*Lath. ind. orn.* p. 280.

---

Africæ incola esse creditur magna hæc avis,  
quam primum descripsisse opinor Lathamum. Pu-  
tatur etiam interdum in India generari. Iconem  
pulcherrimam mutuati sumus a tabulis miscellaneis  
quas pereleganter edidit Dominus Millerus.









THE  
RED-BILLED HOOPOE.



*GENERIC CHARACTER.*

*Bill* bowed, convex, somewhat compressed, rather obtuse.

*Tongue* obtuse, perfectly entire, triquetrous, very short.

*Feet* formed for walking.

*SPECIFIC CHARACTER, &c.*

Violet-black HOOPOE, with a gloss of green; the tail-feathers marked on each side towards the tips with a spot of white; the bill red.


Red-billed Promerops.

*Lath. Synops. Suppl. 1. p. 124.*

*Millar. Illustr. nat. hist. pl. 52.*

---

This magnificent bird, which seems to have been first described by Mr. Latham, is supposed to be a native of Africa; but it is also said to be found in India. The figure here given is copied from the beautiful representation published by Mr. Millar in his splendid plates of natural history.



# SPHAGEBRANCHUS ROSTRATUS.



## *CHARACTER GENERICUS.*

Spiracula duo sub collo.

*Bloch. ichth. 12. p. 78.*

## *CHARACTER SPECIFICUS.*

**SPHAGEBRANCHUS** capite rostrato.

*Bloch. ichth. 12. p. 79.*

APODES.


---

Genus Sphagebranchus, a Blochio primum institutum, hanc unicam continet speciem, in mari Indico præcipue repertam, et in longitudinem forte Anguillæ vulgaris crescentem.





R9X



THE  
ROSTRATED SPHAGEBRANCHUS.



*GENERIC CHARACTER.*

*Spiracula* two, situated beneath the neck.


*SPECIFIC CHARACTER.*

SPHAGEBRANCHUS with the head terminating in a sharp-pointed snout.

---

The genus *Sphagebranchus*, instituted by Dr. Bloch, contains only the species represented on the present plate. It is chiefly seen in the Indian seas, and is supposed to arrive at the size of a common eel.





## BOA ANTARCTICA.



### *CHARACTER GENERICUS.*

*Scuta* abdominalia et subcaudalia.

### *CHARACTER SPECIFICUS, &c.*

BOA gilva, fasciis creberrimis transversis rufis,  
subtus flavescens, nigro maculata.

*Scut: abdom.* 113.

*subcaud.* 27.

*Squam: terminales* 22.

---

Venenata admodum est hæc species; rara insuper, nec antea descripta. Longitudo ut plurimum pedalis pertingit interdum ad quindecim uncias. Insignis est crassitudo corporis. Ob colores notabiles maculasque ex æquo sitas vix poteris permiscere hanc speciem cum ulla alia Boa quam adhuc novimus. In Australasia generatur Boa Antarctica.







272

335

THE  
ANTARCTIC BOA.



*GENERIC CHARACTER.*

*Scuta* or undivided plates both on the abdomen  
and beneath the tail.

*SPECIFIC CHARACTER, &c.*

Gilvous BOA, with numerous transverse rufous  
fasciæ; beneath yellowish, spotted with  
black.


*Abdominal plates* 113.

*Subcaudal* 27.

*Terminal scales* 22 pair.

---

This beautiful, and hitherto undescribed serpent, is a native of Australasia, and is considered as a highly poisonous species. Its general length is about twelve or fifteen inches: its form remarkably thick in proportion. From the singularity of its colours and the regularity of its markings it is not easily confounded with any other species.



# SCARABÆUS LONGIMANUS.



## CHARACTER GENERICUS.

*Antennæ* clavatæ capitulo fissili.

*Tibiæ* anticæ sæpius dentatæ.

*Lin. Syst. Nat.* p. 541.

## CHARACTER SPECIFICUS, &c.

SCARABÆUS muticus, pedibus anticis arcuatis longissimis.

*Lin. Syst. Nat.* p. 549.


*Fabr. spec. ins.* p. 18.

*Voet. Scarab.* t. 11. f. 97.

---

In India innascitur rarissima hæc species, cujus veram magnitudinem cernere est in tabula.









282

282

THE  
LONG-ARMED BEETLE.



*GENERIC CHARACTER.*

*Antennæ* divided at the tip or head into several lamellæ.

*Tibiæ* or second joints of the fore-legs generally toothed.

*SPECIFIC CHARACTER, &c.*

Smooth chesnut-brown BEETLE, with very long, crooked fore-legs.


The crook-legged Indian BEETLE.

La Girafette.

*Pl. Enl. 41. f. 1.*

---

This highly rare insect is a native of India, and is represented on the plate in its natural size.



## TANAGRA FESTIVA.



### CHARACTER GENERICUS.

*Rostrum* conicum, acuminatum, emarginatum,  
basi subtrigonum, apice declive.

*Lin. Syst. Nat.* p. 313.

### CHARACTER SPECIFICUS, &c.

TANAGRA viridis, capite cœruleo, collari(maris)  
rubro, humeris nigris, flavo marginatis.

TANAGRA Tricolor. Var.  $\beta$ .

*Lath. ind. orn.* p. 428.

TANAGRA Cayanensis varia cyanocephalos.

*Briss. Suppl.* p. 62. t. 4. f. 2.

---

Formosior est Tanagra festiva seu rubricollis vel ipsa Tanagra *Tatao*, quam in hoc opere jam descripsimus. Cayennam inhabitat; et cernere est in tabula naturalem avis magnitudinem. Collari isto splendide sanguineo quo mas ornatur caret femina.

1. The first part of the document discusses the importance of maintaining accurate records of all transactions and activities. It emphasizes that this is crucial for ensuring transparency and accountability in the organization's operations.

2. The second part of the document outlines the various methods and tools used to collect and analyze data. It highlights the need for consistent and reliable data collection processes to support informed decision-making.

3. The third part of the document focuses on the role of technology in modern data management. It discusses how advanced software solutions can streamline data collection, storage, and analysis, thereby improving efficiency and accuracy.

4. The fourth part of the document addresses the challenges associated with data security and privacy. It stresses the importance of implementing robust security measures to protect sensitive information from unauthorized access and breaches.

5. The fifth part of the document explores the ethical implications of data collection and analysis. It discusses the need for transparency in data practices and the importance of respecting individual privacy and consent.

6. The sixth part of the document provides a summary of the key findings and recommendations. It reiterates the importance of a data-driven approach and offers practical advice for organizations looking to optimize their data management processes.







THE  
COLLARED TANAGER.



*GENERIC CHARACTER.*

*Bill* conical, acuminate, a little inclining towards the point; upper mandible slightly ridged, and notched near the end.

*SPECIFIC CHARACTER, &c.*

Green TANAGER, with blue head, red collar (in the male) and black shoulders edged with yellow.

The Red-collared TANAGER.

Le Tricolor.


*Buff.* 4. p. 276.

Green-headed TANAGER. Var. A.

*Lath. Syn.* 2. p. 235.

---

This bird, which perhaps exhibits a more brilliant plumage than even the *Tanagra Tatao* described in a former number of the present work, is a native of Cayenne, and is represented in its natural size on the annexed plate. The female is destitute of the red collar which forms so conspicuous an ornament in the male.



# CATAPHRACTUS COSTATUS.



## CHARACTER GENERICUS.

*Corpus* Cataphractum.

*Os* terminale. *Bloch. ichth.* 11. p. 65.

*Abdominales.*

## CHARACTER SPECIFICUS, &c.

CATAPHRACTUS ordine scutorum simplici,  
pinna caudæ lunata.

*Bloch. ichth.* 11. p. 66.

*Mystus cirris* sex longissimis, &c.

*Gronov. Mus.* 2. p. 24.

Silurus CATAPHRACTUS. S. pinna dorsali  
postica adiposa squamis serie simplici, cirris  
sex, cauda bifida.


*Lin. Syst. Nat.* p. 506.

---

A Blochio institutum genus *Cataphractus* perpaucas continet species, quæ ut in Siluris disponentur auctor erat Linnæus. Cataphractum costatum, qui exemplum præbeat generis, selegimus, maria incolentem Americana, et corpus habentem præduris laminis armatum. Magnitudo ei ut plurimum est quasi *Cyprini Carpionis*.







THE  
RIBBED SOLDIER-FISH.



*GENERIC CHARACTER.*

*Body* mailed.

*Mouth* terminal.

*SPECIFIC CHARACTER, &c.*

**SOLDIER-FISH** with a simple row of shields,  
and lunated tail.

The American **SOLDIER-FISH**.

The **RIBBED SILURE**.

---

The genus *Cataphractus*, instituted by Dr. Bloch, contains but a very small number of species, which by Linnæus were arranged under the genus *Silurus*. The *Cataphractus costatus*, selected as an example of the genus, is a native of the American seas, and is remarkable for the very strong or bony armature of the body. It is about the size of a common Carp.





## ACTINIA DIANTHUS.



### CHARACTER GENERICUS.

*Corpus* se affigens basi, carnosum, oblongum, teres, contractile, viviparum.

*Os* terminale, dilatabile, tentaculis cinctum.

*Apertura* præter os nulla.

### CHARACTER SPECIFICUS, &c.

ACTINIA lævis subcylindrica, disco quinquepartito foliaceo, tentaculis exiguis albis ornato, osculo elevato striato.


*Ellis. Zooph. p. 7. Act. Angl. vol. 57.*  
*p. 436. t. 19. f. 6. 7.*

Hydra DIANTHUS.

*Lin. Syst. Nat. Gmel. p. 3869.*

---


Formosam hanc Actiniam in Britannia innasci docet qui primus eam descripsisse videtur celeberrimus Ellisius, rupesque oræ Sussexiensi adjacentes sæpius incolere, eas præcipue quæ Hastingsarum ex adverso sitæ sunt. Eadem est magnitudine quam ostendit tabula, et eodem fere utitur vivendi modo quo reliquum genus.





150

539



R.P.V.

THE  
CARNATION ACTINIA.



*GENERIC CHARACTER.*

*Body* fixing itself by the base, fleshy, oblong, cylindrical, contractile, viviparous.

*Mouth* terminal, expansile, surrounded with tentacula.

No other opening but the mouth.

*SPECIFIC CHARACTER, &c.*


Smooth subcylindric Actinia, with foliaceous five-parted disk beset with minute tentacula, and elevated, striated mouth.

Sea CARNATION.

*Ellis. Zooph. p. 7. Phil. Trans. vol. 57.*  
p. 436. pl. 19. f. 8.

---

The beautiful species of Actinia represented on the present plate seems to have been first described by Mr. Ellis, who informs us that it is an native of our own island, and is principally found on the rocks of the Sussex coast, and more especially on those which are opposite the town of Hastings. In its general properties and manners it resembles the rest of its tribe, and is expressed on the plate in its natural size.





1





540

540

## PAPILIO DEÏPHOBUS.



### CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatae.

*Alæ* sedentis erectæ sursumque conniventes, (volatu diurno.)

*Lin. Syst. Nat.* p. 744.

### CHARACTER SPECIFICUS, &c.

PAPILIO alis caudatis fuscis, superioribus macula humerali rubra, posticis apice fulvis nigro maculatis.

PAPILIO DEÏPHOBUS? P. alis caudatis nigris, subtus basi rubro maculatis, posticis maculis septem rubris subannularibus.

*Lin. Syst. Nat.* p. 746.

*Eq. Tr.*

PAPILIO Alcander?

*Cram.* 1. p. 64. t. XL.

---

Indiam incolit formosum hoc insectum, cujus veram magnitudinem ostendit tabula. Coloribus interdum variat.

# DEIPHOBUS.



## GENERIC CHARACTER.

*Antennæ* thickening towards the upper part,  
and generally terminating in a knob.

*Wings* (when sitting) erect, and meeting upwards.

(Flight diurnal.)

## SPECIFIC CHARACTER, &c.

Caudated brown Butterfly, with the upper wings marked by a red shoulder-spot; the lower with fulvous tips spotted with black.

Great dusky Swallow-tailed Butterfly.

*Edw. av. pl. 347.*

---

This beautiful Insect is a native of India, and is represented in the plate in its natural size. The different specimens are observed to vary occasionally in their colours.

## INDEX.

- PL.
503. *Anableps tetropthalmus*.  
 507. *Aplysia depilans*.  
 525. *Alcedo chlorocephala*.  
 532. *Ascidia Aurantium*.  
 539. *Actinia Dianthus*.  
 535. *Boa antarctica*.  
 524. *Cancer cristatus*.  
 496. ——— *Phalangium*.
538. *Cataphractus costatus*.  
 495. *Conus Ammiralis* et var.  
 502. ——— *arausiacus* et  
*arachr*.  
 523. ——— *Tulipa*.  
 509. *Coracias Sinensis*.
506. *Coryphæna cærulea*.  
 511. *Echeneis Remora*.  
 526. *Eques Americanus*.  
 505. *Fringilla elegans*.  
 518. *Fistularia Tabacaria*.  
 514. *Holocentrus tigrinus*.  
 522. ——— *Sogo*.  
 501. *Lemur Calago*.  
 521. *Lanius bicolor*.  
 527. *Murex ramofus*.
515. *Nautilus Pompilius*.
500. *Papilio Agenor*.  
 504. ——— *Protesilaus*.  
 512. ——— *Æneas & Peran-*  
*thus*.  
 516. ——— *Panthous*.  
 540. *Deïphobus*.  
 529. *Pelecanus Sinensis*.  
 520. *Phalæna macroura*.  
 530. 531. — *Agrippina*.  
 493. *Picus pileatus*.  
 497. ——— *principalis*.  
 498. *Scomber Ductor*.  
 494. ——— *ruber*.

## INDEX.

- PL.
503. *Anableps four-eyed*.  
 539. *Actinia carnation*.  
 532. *Ascidia orange*.  
 507. *Aplysia Mediterranean*.  
 536. *Beetle long-armed*.  
 535. *Boa antarctic*.  
 500. *Butterfly Agenor*.  
 512. ——— *Æneas and Pe-*  
*ranthus*.  
 516. ——— *Panthous*.  
 504. ——— *Protesilaus*.  
 540. ——— *Deïphobus*.
495. *Cone Admiral* and var.  
 502. ——— *Orange - flag* and  
*Cobweb*.  
 523. ——— *Tulip*.  
 506. *Coryphæna blue*.  
 496. *Crab slender-legged*  
 524. ——— *crested*.  
 505. *Finch variegated*.  
 518. *Fistularia fibre-tailed*.  
 522. *Holocentrus red*.  
 514. ——— *variegated*.  
 533. *Hoopoe red-billed*.  
 513. *Humming-Bird topaz-*  
*throated*.  
 517. ——— *stripe -*  
*checked*.  
 525. *Kingfisher green-headed*  
 526. *Knight-fish American*.  
 501. *Lemur Calago*.
494. *Mackrel red*.  
 498. *Mackrel Pilot*.  
 527. *Murex branched*.  
 515. *Nautilus great*.  
 529. *Pelican Chinese*.  
 520. *Phalæna long-tailed*.  
 529. ——— *Agrippina*.  
 504. *Roller Chinese*.  
 511. *Remora fork-tailed*.

# I N D E X.

- | Pl.                                   | Pl.                                  |
|---------------------------------------|--------------------------------------|
| 536. <i>Scarabæus longimanus.</i>     | 521. Shrike blue.                    |
| 499. <i>Strombus Fusus.</i>           | 538. Soldier-fish ribbed.            |
| 519. ----- <i>latissimus.</i>         | 534. <i>Sphagebranchus rostrated</i> |
| 534. <i>Sphagebranchus rostratus.</i> | 528. Sphinx Jasmine.                 |
| 528. <i>Sphinx Atropos.</i>           | 499. <i>Strombus Spindle.</i>        |
| 510. <i>Tethys Fimbria.</i>           | 519. ----- <i>Wing.</i>              |
| 537. <i>Tanagra festiva.</i>          | 510. <i>Tethys crenulated.</i>       |
| 513. <i>Trochilus Pella.</i>          | 537. <i>Tanager collared.</i>        |
| 517. ----- <i>superbus.</i>           | 508. <i>Volute Ethiopian.</i>        |
| 508. <i>Voluta Æthiopica.</i>         | 493. <i>Woodpecker pileated.</i>     |
| 533. <i>Upupa erythrorynchus.</i>     | 497. ----- <i>White -</i>            |
|                                       | billed.                              |











